

09/462416



STATE OF ISRAEL

PCT/IL 98 / 00321

REC D 23 JUL 1998

WIPO PCT

This is to certify that
annexed hereto is a true
copy of the documents as
originally deposited with
the patent application
particulars of which are
specified on the first page
of the annex.

זאת לתעודה כי
רצופים כזה העתקים
נכונים של המסמכים
שהופקרו לכתחילה
עם הבקשה לפטנט
לפי הפרטים הרשומים
בעמוד הראשון של
הנספח.

PRIORITY DOCUMENT

This 16-07-1998 היום

לשם
הפטנטים

רשם הפטנטים
Registrar of Patents

נתאשר

לשימוש הלשכה
For Office Use

חוק הפטנטים, תשכ"ז - 1967
PATENT LAW, 5727 - 1967

ב ק ש ה ל פ ט נ ט
Application for Patent

121284	מספר: Number
10-07-1997	תאריך: Date
	הוקדם/נדרחא Ante/Post-dated

אני, (שם המבקש, מענו ולגבי גוף מאוגד - מקום התאגדותו)
I (Name and address of applicant, and in case of body corporate-place of incorporation)

Yeda Research and Development Co. Ltd.,
Israeli Company,
at the Weizmann Institute of Science,
P.O. Box 95,
Rehovot 76100, Israel

ידע חברה למחקר ופיתוח בע"מ,
חברה ישראלית,
ליד מכון וייצמן למדע,
ת.ד. 95
רחובות 76100


Assignment העברה
שם ההמצאה הוא בעל אמצאה מכח
of an invention the title of which is Owner, by virtue of

IL-6 - ובין ה - IL-6
חלבון אחוי בין הקולטן הממס ל - IL-6
אנלוגים של חלבון האחוי ושימושיהם (בעברית)
(Hebrew)

Chimeric Interleukin-6 Soluble Receptor/Ligand Protein, Analogs
thereof and Uses thereof (באנגלית)
(English)

hereby apply for a patent to be granted to me in respect thereof.

מבקש כואת כי ינתן לי עליה פטנט

• בקשת חלוקה • Application of Division		• בקשת פטנט מוסף • Application for Patent Addition		• דרישה רין קדימה • Priority Claim		
מבקשת פטנט from Application	לבקשה/לפטנט to Patent/Appl.	מספר/סימן Number/Mark	תאריך Date	מדינת האגוד Convention Country		
No. dated	No. dated					
• יסוי כת: כללי / מיוחד - רצוף בזה / עוד יוגש O.A.: general/individual-attached/to be filed later- הוגש בענין..... המקן למסירת מסמכים בישראל Y/97-44 Address for Service in Israel הנרי עינב, עורך פטנטים אגנטר-לאב בע"מ, קרית ויצמן נמ-צגונה 76110						
חתימת המבקש Signature of Applicant		היום This				
For the Applicant :  Henry Einav		10 בחודש יולי שנת 1997 10 of the year July of 1997				
		לשימוש הלשכה For Office Use				

CHIMERIC INTERLEUKIN-6 SOLUBLE RECEPTOR/LIGAND
PROTEIN, ANALOGS THEREOF AND USES THEREOF

חלבון אחוי בין הקולטן הממיס ל - IL-6
ובין ה- IL-6, אנלוגים של חלבון האחוי ושימושיהם

Yeda Research and Development Co. Ltd.

Y/97-44

Inventors : Michel Revel, Judith Chebath, Tsvee Lapidot, Orit Kollet

Field of the invention

The present invention is generally in the field of interleukin-6 (IL-6) biological activities which are dependent on the agonistic action of soluble IL-6 receptor (sIL-6R). More specifically, the present invention concerns a novel chimeric sIL-6R/IL-6 protein constructed from the fusion of essentially the naturally occurring form of sIL-6R and IL-6, and biologically active analogs thereof, which are particularly useful for treating cancer, via inhibition of cancerous cell growth and for enhancing bone marrow transplantation.

Background of the Invention and Prior Art

Interleukin-6 (IL-6) is a well known cytokine whose biological activities are mediated by a membranal receptor system comprising two different proteins one named IL-6 Receptor (IL-6R or gp80) and the other gp130 (reviewed by Hirano et al, 1994). Soluble forms of IL-6R (sIL-6R), corresponding to the extracellular domain of gp80, are natural products of the human body found as glycoproteins in blood and in urine (Novick et al, 1990, 1992). An exceptional property of sIL-6R molecules is that they act as potent agonists of IL-6 on many cell types including human cells (Taga et al, 1989; Novick et al, 1992). This is due to the fact that even without the intracytoplasmic domain of gp80, sIL-6R is still capable of triggering the dimerization of gp130 in response to IL-6, which in turn mediates the subsequent IL-6-specific signal transduction and biological effects (Murakami et al, 1993). The active IL-6 receptor complex is in fact a hexameric structure formed by two gp130 chains, two IL-6R and two IL-6 ligands (Ward et al, 1994; Paonessa et al, 1995), in which sIL-6R has two types of interaction with gp130 both of which are essential for the IL-6-specific biological activities (Halimi et al, 1995).

Treatment with sIL-6R results in an enhancement of the biological activities of IL-6 in many cell types. An example is tumor cells whose growth is inhibited to a greater extent by IL-6 when sIL-6R is added, such as murine myeloleukemic M1

cells (Taga et al, 1989), human breast carcinoma T47D cells (Novick et al, 1992) or human Non-small cell lung carcinoma cells (Ganapathi et al, 1996). While IL-6 has anti-metastatic activities *in vivo* (Katz et al, 1995), sIL-6R can, however, also enhance such *in vivo* anti-tumor effects of IL-6 (Mackiewicz et al 1995). Another activity of IL-6 which is enhanced by sIL-6R addition is the stimulation of hematopoietic stem cells to produce multilineage colonies (Sui et al, 1995). The present inventors have also observed that the survival of primary cultures of brain oligodendrocytes is supported by the sIL-6R and IL-6 combination (Oh, 1997), while, however, IL-6 alone is poorly active in such cultures (Kahn and De Vellis, 1994). This finding indicates that IL-6 when combined with sIL-6R can mimick the activity of other neurotropic cytokines such as Ciliary Neurotropic Factor (CNTF) or Leukemia Inhibitory Factor (LIF) which also act through gp130, as is also the case for IL-11 and Oncostatin M (Hirano et al, 1994).

In an attempt to provide a molecule which may combine the above noted functions of IL-6 and sIL-6R, there has recently been reported the production in recombinant yeast cells of a fusion protein between a truncated segment of the human IL-6R sequence and IL-6, linked by a glycine-rich linker (Fischer et al., 1997). This fusion protein includes essentially only the IL-6R cytokine receptor N-domain and the cytokine receptor C-domain, and thus lacks essentially all of the IL-6R immunoglobulin (Ig)-like domain, and the receptor pre-membrane region, and as such represents even a truncated form of the sIL-6R, this truncated sIL-6R in the fusion protein being linked via the above noted glycine-rich linker to essentially the whole mature form of IL-6. Besides lacking parts of the natural sIL-6R, this fusion protein by being produced in yeast cells, most likely does not have the glycosylation pattern that such a fusion protein would have if it were produced in mammalian cells, in particular, in human cells. In fact, this yeast produced fusion protein has a molecular weight of only about 57 kDa in contrast to a fusion product containing essentially all of the natural sIL-6R and IL-6 amino acid residues and

being fully glycosylated in mammalian (e.g. human) cells, which has a molecular weight of about 85 kDa (see Example 2 herein below).

The common experience in developing recombinant proteins which can be used for treating human patients has shown that it is important to remain as close as possible to the natural forms of the proteins, as they are found in the human body, in order to avoid triggering of antibodies and other side effects observed with non-natural recombinant products. For this reason, it has been advantageous to use recombinant mammalian cell systems to produce glycosylated proteins such as Interferon- β or Granulocyte-colony stimulating factor (Chernajovsky et al, 1984, Holloway, 1994) in a chemical form as similar as possible to the natural human product. Bacteria or microorganisms such as, for example, yeasts, which do not glycosylate properly, also cause the wrong folding of the protein chains, leading to immunogenic reactions. This is particularly important in respect of IL-6 which is heavily modified postrationally by N- and O- glycosylation as well as by phosphorylation (Revel, 1989 for review), and in respect of the natural sIL-6R from human blood and urine which is a glycoprotein whose N-terminus and C-terminal amino-acids are constant and have been determined (Novick et al, 1990 and co-owned patents by the present inventors Nos. US 5,216,128 and its corresponding EP 413908 B1).

Accordingly, it would seem that the above noted previous fusion product between part of the sIL-6R and IL-6 has a number of possible drawbacks especially as regards its use for treating humans and this in view of its lacking part of the sIL-6R and its production in yeasts which may provide for incorrect glycosylation of the protein.

Heretofore, a fusion molecule comprising the natural sIL-6R found in human body fluids and the natural IL-6, and which is produced in human or other mammalian cells, has not been described.

It is therefore an aim of the present invention to provide such a fusion molecule comprising the natural sIL-6R and the natural IL-6 which is produced in mammalian cells.

It is another aim of the present invention to use such a fusion protein (sIL-6R/IL-6 chimera) to inhibit the growth of highly metastatic melanoma cells at very low concentrations, these cells being resistant to IL-6 or sIL-6R alone.

Yet another aim is to use such a fusion protein (the sIL6R/IL-6 chimera) for the *in vivo* engraftment of human hematopoietic stem cells in bone marrow transplantation protocols.

A further aim of the invention is to provide pharmaceutical compositions which contain the above mentioned natural sIL-6R-natural IL-6 fusion protein (sIL-6R/IL-6 chimera) for the treatment of cancer and for use in bone marrow transplantation procedures.

Other aims and aspects of the present invention will be set forth or will arise from the following disclosure of the present invention.

Summary of the Invention

In accordance with the present invention there have been produced a number of fusion proteins (chimeras) each comprising essentially all of the naturally occurring sIL-6R from human body fluids and essentially all of the mature form of the naturally occurring human IL-6, and each joined by short linker peptides which can be as short as 3 amino acid residues in length or longer, for example, 13 amino acid residues in length (see below and Examples 1 and 2). It should be noted, however, that in these fusion proteins the linker peptides may be omitted and the sIL-6R moiety may be directly linked to the IL-6 moiety. Since linkers representing non-natural amino acid sequences may be immunogenic epitopes eliciting antibodies in patients, it is preferable to have a directly fused sIL-6R/IL-6 chimera that has the desired biological activity, while at the same time there is minimized

the risk of inducing such potentially deleterious antibody formation when such a chimera is administered.

The conservation of the entire sIL-6R sequence including the Ig-like domain as found in the naturally occurring molecule, as well as the proper glycosylation and other post-translational modification introduced by human or mammalian cells when the above chimera is produced in such cells, are also important to reduce the potential immunogenicity of the chimeric protein product.

However, it is possible to use a very short linker of above three-four amino acids at the fusion point between the sIL-6R and IL-6 moieties of the chimeric protein. Such a short linker would serve to separate these two moieties but would not be an immunogenic epitope. It is of course also possible to use longer linkers of up to about 30 amino acids to provide for separation between the two moieties but here care must be taken and biological efficacy and safety experiments must be performed to ensure that chimeric molecules with such linkers are not immunogenic.

In fact, it has been surprisingly shown in accordance with the present invention that such longer linkers are not essential for the activity of the chimeric protein indicating that proper folding of the chimera does not require a longer linker especially when essentially all of the naturally-occurring sequences of the sIL-6R and IL-6 moieties are incorporated into the chimeric molecule (see Example 3 and Fig. 5 which relate also to a comparison between an sIL-6R/IL-6 chimera having a very short (3 amino acids) linker and a similar chimera having a longer linker of 30 amino acids).

These fusion proteins or sIL-6R/IL-6 chimeras have been efficiently produced, in accordance with the present invention, in mammalian cell expression systems to yield glycosylated products having potent activity on tumor cells which are usually non-responsive to IL-6 or sIL-6R alone, and which were highly effective on ensuring the success of engraftment of human bone marrow transplanted cells

(see below and Examples 1-4). In fact, in such bone marrow transplants, the sIL-6R/IL-6 chimeras were essential for the survival and proliferation of the transplanted non-committed pluripotential hematopoietic stem cells. Moreover, from the experimental results presented herein below as well as other analyses it arises that various analogs of the sIL-6R/IL-6 chimeric protein of the invention can be prepared which have essentially the same biological activity of the sIL-6R/IL-6 chimera, these analogs being sIL-6R/IL-6 chimeras in which one or more amino acid residues have been deleted, added or substituted by others, the only limitation on such analogs being that they retain most of the naturally occurring sIL-6R and IL-6 sequence. For example, amino acid additions to the naturally occurring sIL-6R and IL-6 sequences are preferably limited to up to between about 20 amino acids, and preferably these additions are at the site of junction between the sIL-6R and IL-6, i.e. the linker molecule. Likewise, deletions from the sIL-6R and IL-6 sequences are preferably limited to up to between about 20-30 amino acids; and substitutions of amino acid residues in the sIL-6R and IL-6 sequences by other amino acid residues are preferably also limited to up to between about 20-30 amino acids. All of the aforesaid deletions, additions and substitutions are acceptable in accordance with the present invention when the so-modified analogs that are obtained retain essentially the biological activity of the sIL-6R/IL-6 chimera composed of essentially the naturally-occurring sequences, and retain essentially the same glycosylation pattern of the chimera composed of essentially the naturally-occurring sequences when expressed in mammalian cells.

Accordingly, the present invention provides a chimeric glycosylated soluble interleukin-6 receptor (sIL-6R)-interleukin-6 (IL-6) protein (sIL-6R/IL-6) and biologically active analogs thereof, comprising a fusion protein product between essentially all of the naturally occurring form of sIL-6R and essentially all of the naturally occurring form of IL-6, said sIL-6R/IL-6 and analogs thereof being

glycosylated in a similar fashion to the glycosylation of naturally occurring sIL-6R and IL-6.

Embodiments of the above chimeric protein of the invention include :

(i) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, wherein said sIL-6R is fused to IL-6 via a peptide linker molecule.

(ii) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as in (i) above, wherein said linker is a very short, non-immunogenic linker of about 3-4 amino acid residues.

(iii) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as in (ii) above, wherein said linker is a tripeptide of the sequence E-F-M (Glu-Phe-Met).

(iv) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as in (i) above, wherein said linker is a peptide of 13 amino acid residues of sequence E-F-G-A-G-L-V-L-G-G-Q-F-M (Glu-Phe-Gly-Ala-Gly-Leu-Val-Leu-Gly-Gly-Gln-Phe-Met).

(v) A chimeric sIL-6R/IL-6 protein, being the herein designated sIL6R δ Val/IL6 having a tripeptide linker of sequence E-F-M between the C-terminal Val-356 of sIL-6R and the N-terminal Pro-29 of IL-6R, said chimeric protein having the sequence set forth in Fig. 3.

(vi) A chimeric sIL-6R/IL-6 protein, being the herein designated sIL6R δ Val/L/IL6 having a 13 amino acid peptide linker of sequence E-F-G-A-G-L-V-L-G-G-Q-F-M between the C-terminal Val-356 of sIL-6R and the N-terminal Pro-29 of IL-6R, said chimeric protein having the sequence set forth in Fig. 3 wherein the tripeptide of sequence E-F-M between positions 357-359 of Fig. 3 is replaced by said 13 amino acid peptide sequence.

(vii) A chimeric sIL-6R/IL-6 protein, wherein said protein is produced in mammalian cells in a fully processed form.

(viii) A chimeric sIL-6R/IL-6 protein, wherein said protein is produced in human cells.

(ix) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as above, wherein said chimeric protein and analogs are characterized by being capable of inhibiting the growth of highly malignant cancer cells.

(x) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as above, wherein said chimeric protein and analogs are characterized by being capable of inhibiting the growth of highly malignant melanoma cells.

(xi) A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, as above, wherein said chimeric protein and analogs are characterized by being capable of eliciting the *in vivo* engraftment of human hematopoietic cells in bone marrow transplantations.

The present invention also provides a DNA sequence encoding a chimeric sIL-6R/IL-6 protein and biologically active analogs thereof as noted above according to the invention.

In addition, the present invention also provides a DNA vector comprising a DNA sequence encoding a chimeric sIL-6R/IL-6 protein and biologically active analogs thereof of the invention, as noted above, said vector being suitable for expression of said chimeric protein in mammalian cells.

Embodiments of the DNA vector of the invention include :

(i) A DNA vector wherein said vector is suitable for expression of said chimeric protein in human cells.

(ii) A DNA vector wherein when said vector is expressed in mammalian or human cells, the expressed chimeric protein has a sequence that permits full processing of the chimeric protein by the mammalian or human cell and secretion of the fully processed chimeric protein from the cells into the culture medium in which said cells are grown.

(iii) A DNA vector, as above, wherein said vector is the herein designated plasmid pcDNA_{sIL-6R/IL-6} comprising a pcDNA3 vector containing the DNA sequence encoding the chimeric sIL-6R/IL-6 protein under the control of a cytomegalovirus (CMV) promoter.

(iv) A DNA vector, as above, wherein said vector is the herein designated plasmid pcDNA sIL-6R/L/IL-6 comprising a pcDNA3 vector containing the DNA sequence encoding the chimeric sIL-6R/IL-6 protein under the control of a cytomegalovirus (CMV) promoter, and wherein in said DNA sequence encoding said chimeric sIL-6R/IL-6 protein there is inserted a linker sequence encoding a linker peptide at the EcoRI site placed between the sequence encoding the sIL-6R part and the sequence encoding the IL-6 part of the protein.

Likewise, the present invention also provides transformed mammalian cells containing a DNA vector as above, that is capable of expressing the sIL-6R/IL-6 chimeric protein sequence carried by said vector and of fully processing the expressed protein and secreting it into the culture medium in which said cells are grown.

An embodiment of these transformed cells are the herein described human embryonal kidney cells 293 (HEK293) transfected by the pcDNA sIL-6R/IL-6 vector, said cells being capable of expressing the sIL-6R/IL-6 chimeric protein, fully processing said protein and secreting said protein into the culture medium in which said cells are grown in the form of an about 85 kDa glycoprotein.

The present invention also provides a method for producing a chimeric protein or biologically active analogs thereof, as above, comprising growing the aforesaid transformed cells under conditions suitable for expression, processing and secretion of said protein or analogs into the culture medium in which said cells are grown; and purifying said protein or analogs from said culture medium by immunoaffinity chromatography using monoclonal antibodies specific for sIL-6R.

The chimeric protein of the present invention has a number of uses including:

(i) use of a chimeric sIL-6R/IL-6 protein or analogs, salts of any one thereof, and mixtures thereof, as an inhibitor of cancer cells.

(ii) use, as in (i) above, as an inhibitor of highly malignant melanoma cells.

(iii) use of a chimeric sIL-6R/IL-6 protein or analogs, salts of any one thereof, and mixtures thereof, as an active ingredient for eliciting engraftment of human hematopoietic cells in bone marrow transplantation.

(iv) use of a chimeric sIL-6R/IL-6 protein or analogs, salts of any one thereof, and mixtures thereof, as an active ingredient for increasing hematopoiesis, for treating neurological conditions, or for other applications in which IL-6 or sIL-6R are used.

Similarly, the chimeric protein of the present invention may be used to prepare medicaments for a number of medical indications, namely, a chimeric sIL-6R/IL-6 protein or analogs, salts of any one thereof and mixtures thereof, for use in the preparation of a medicament for treating cancers by way of inhibition of cancer cells, or in the preparation of a medicament for enhancement of bone marrow transplantation by way of eliciting engraftment of human hematopoietic cells in bone marrow transplantation, or in the preparation of a medicament for increasing hematopoiesis, or in the preparation of a medicament for treating neurological disorders, or in the preparation of a medicament for other applications in which IL-6 or sIL-6R are used.

Moreover, the present invention also provides a pharmaceutical composition comprising as active ingredient a chimeric sIL-6R/IL-6 protein or analog thereof as above, and a pharmaceutically acceptable carrier, diluent or excipient.

Embodiments of this pharmaceutical composition of the invention include :

(i) A pharmaceutical composition for the treatment of mammalian cancers.

(ii) A pharmaceutical composition for the enhancement of bone marrow transplantation.

(iii) A pharmaceutical composition for the treatment of neurological disorders, or for increasing hematopoiesis or for other applications in which IL-6 or sIL-6R are used.

The present invention also provides for a method for treating cancers in mammals, or for enhancing bone marrow transplantations, or for treating neurological disorders, or for increasing hematopoiesis, or for other applications in which IL-6 or sIL-6R are used, comprising administering to a patient a pharmaceutical composition, as above, in a suitable dosage form and by a suitable route of administration.

Other aspects and embodiments of the present invention are set forth or arise directly from the following detailed disclosure of the invention.

Brief Description of the Drawings

Figure 1 depicts a schematic representation of the various vectors, reagents and process steps used in the construction of the chimeric DNA molecule encoding a chimeric protein in which is conserved the structure of the natural form of sIL-6R ending at the Val 356 residue followed by the sequence of the natural, mature, processed form of IL-6, as detailed in Example 1;

Figure 2 (A,B) shows the results obtained from the analysis performed to identify the sIL-6R δ Val/IL-6 p86 chimera by polyacrylamide gel electrophoresis (A) and by bioactivity profile (B), wherein in Fig. 2A there is shown a reproduction of a Coomassie stained gel on which were electrophoresed immunopurified fractions eluted from affinity chromatography columns loaded with a secreted protein sample obtained from cell cultures transfected with a vector encoding the chimeric protein; and in Fig. 2B there is shown a graphic representation of the

biological activity (growth inhibition of F10.9 melanoma cells) of each of the above noted fractions eluted from the affinity chromatography columns, all as detailed in Examples 2 and 3;

Figure 3 depicts schematically the amino acid sequence (one-letter code) of the sIL-6R δ Val/IL-6 chimera in which is shown the different domains of the molecule, including the N-terminal signal peptide (underlined above), the immunoglobulin-like (Ig-like) domain, the cytokine receptor N-domain (underlined below), the cytokine C-domain (underlined above) and the receptor pre-membrane region, all of the sIL-6R part of the chimera; as well as the mature IL-6 moiety (underlined below) of the chimera, as described in Examples 1 and 2;

Figure 4 (A, B) shows photographs of F10.9 melanoma cells in culture without (A) and with (B) treatment with the sIL-6R/IL-6 chimeric protein for 4 days, wherein in Fig. 4B there is apparent the morphological changes induced in such metastatic melanoma cells (F10.9 cells) by treatment with the sIL-6R/IL-6 chimera, as described in Example 3;

Figure 5 is a graphic representation of the results depicting the growth inhibition of F10.9 melanoma cells by the sIL-6R/IL-6 chimeric protein at various concentrations of the chimera ranging from about 0.12 ng/ml to about 150 ng/ml, as described in Example 3;

Figure 6 is a graphic representation of the results depicting the absence of growth-inhibitory effects on F10.9 melanoma cells of either isolated IL-6 alone (dotted upper curve with open squares) at concentrations ranging from 0-40 ng/ml of IL-6 and sIL-6R alone (point of convergence of all curves on vertical axis where IL-6 concentration is zero); as well as the observed growth inhibitory effects when IL-6 and sIL-6R are added together at various concentrations of each wherein the IL-6 concentration ranges from 10 ng/ml to 40 ng/ml, and sIL-6R added at three concentrations of 100 ng/ml, 200 ng/ml and 400 ng/ml for each IL-6 concentration,

as illustrated in the three lower curves (two dotted curves with open triangles and circles and full curve with closed squares), as described in Example 3; and

Figure 7 is a reproduction of an autoradiogram of a Southern blot showing the requirement of the sIL-6R/IL-6 chimeric protein for successful engraftment of human hematopoietic stem cells during bone marrow transplantation in SCID-NOD mice (two right hand lanes representing the mice which received the sIL-6R/IL-6 chimeric protein in addition to the other necessary factors, SCF, FLT-3, and this in contrast to the three left hand lanes which represent mice having received only SCF and FLT-3 and SCF, FLT-3 as well as isolated, i.e. non-fused, IL-6 and sIL-6R), as described in Example 4.

Detailed description of the invention

The present invention concerns a chimeric sIL-6R/IL-6 protein and biologically active analogs thereof which have essentially all of the naturally occurring form of sIL-6R and essentially all of the naturally occurring form of IL-6 fused together, the site of fusion of which may be by way of a linker peptide, as short as 3 amino acids, and which chimeric sIL-6R/IL-6 protein or analogs have a similar amount and pattern of glycosylation than that of naturally occurring sIL-6R and IL-6. Such a chimeric sIL-6R/IL-6 protein produced in accordance with the present invention in mammalian cells, in particular, in human cells (see Examples 1-4 below) was found to be efficiently expressed in such cells, to be highly glycosylated, and to have potent activity on tumor cells which have no response at all to IL-6 or sIL-6R alone.

More particularly, in accordance with the present invention it has been observed (see Examples 1-3 below) that the aforesaid chimeric sIL-6R/IL-6 protein of the invention causes growth arrest of highly malignant mammalian cells such as the F10.9 melanoma cells at concentrations lower than needed when a mixture of non-fused sIL-6R and IL-6 is used. This is a particularly significant result in view

of the fact that such F10.9 melanoma cells continue to grow normally when treated with only IL-6 or only sIL-6R separately, and undergo growth arrest only when exposed to relatively high dosages of a combination of non-fused IL-6 and sIL-6R. Accordingly, the chimeric sIL-6R/IL-6 protein of the present invention is surprisingly a more potent inhibitor of these highly malignant melanoma cells than a mixture of its separate parts, i.e. a mixture of non-fused IL-6 and sIL-6R. The chimeric protein of the present invention is thus particularly useful as an active ingredient for treating various kinds of cancers.

Furthermore, it has also been found in accordance with the present invention (see Example 4 below) that a chimeric sIL-6R/IL-6 molecule of the present invention is particularly useful for enhancing bone marrow transplantation. In fact, using a known protocol for engraftment of human bone marrow cells into severe combined immunodeficient (SCID) mice, in which stem-cell factor (SCF) and Flt3-ligand are used for enabling survival and proliferation of the most primitive pluripotential hematopoietic stem cells capable of long-term engraftment into recipient bone marrow, it was found that these two factors, SCF and Flt3-ligand, were insufficient to promote the engraftment of human cells into the recipient mouse bone marrow, and that only when the chimeric sIL-6R/IL-6 protein was also added was engraftment successful. This finding indicates that the chimeric protein may be essential in such engraftment protocols. In the same experiments, non-fused IL-6 and sIL-6R when added separately, were insufficient to promote successful bone marrow transplantation and when added together were much less active than the chimeric sIL-6R/IL-6 protein, i.e. at an effective concentration of 100 ng/ml the sIL-6R/IL-6 chimeric protein promoted successful bone marrow transplantation, while the two separate non-fused sIL-6R and IL-6 when added together at even higher concentrations (sIL-6R from 125-1250 ng/ml, IL-6 from 50-200 ng/ml), were much less active in promoting such transplantation.

The above chimeric sIL-6R/IL-6 protein of the invention is preferably a recombinant glycosylated sIL6R/IL6 chimera produced in human cells or in any other suitable mammalian cell expression system which is capable of glycosylating proteins as do human cells and which introduces the same post-translational modifications as do human cells. An important characteristic is that the chimeric glycoprotein so-produced is processed and modified as are the natural sIL-6R and IL-6 parent molecules found in the human body, without truncation and without addition of extraneous unnatural polypeptide sequences, with the exception of the very short tripeptide or when a longer linker peptide is incorporated between the sIL-6R and IL-6 moieties of the chimeric protein.

To prepare the above preferred chimeric protein of the invention, the following features of the naturally-occurring sIL-6R and IL-6 were considered : It is known that the IL-6R present in human cell membranes is produced by a cDNA encoding 468 aminoacids comprising a signal peptide, an Immunoglobulin (Ig) like domain, a cytokine binding domain, a transmembrane region and a cytoplasmic domain (Yamasaki et al, 1988). A soluble form of sIL-6R is found in body fluids which has, like the mature IL-6R from membranes, an N-terminus corresponding to Leu-20 (Novick et al, 1990) and a C-terminus corresponding to Val-356 just before the transmembrane region of IL-6R (see co-owned U.S. Pat. No. 5,216,128 and EP 413.908 B1). In order to fuse this sIL-6R sequence to IL-6, an EcoRI restriction site was introduced following Val-356. The sequence of the mature IL-6 starting at Pro-29 of the IL-6 cDNA and ending at Met-212 (Zilberstein et al, 1986; Hirano et al, 1986) was introduced after this EcoRI site. At this EcoRI site there could also, but not necessarily, be introduced a linker peptide of desired length to distance the sIL-6R and IL-6 moieties from each other in the chimeric protein. As set forth in the Examples below, two different chimeric proteins were produced as examples of such possible chimeric proteins, one having a tripeptide linker and the other having

a 13-amino acid residue linker at this EcoRI site, both being essentially equally active biologically.

The present invention also concerns analogs of the above chimeric sIL-6R/IL-6 protein of the invention, which analogs retain essentially the same biological activity of the chimeric protein having essentially only the naturally occurring sequences of sIL-6R and IL-6. Such analogs may be ones in which up to about 30 amino acid residues may be deleted, added or substituted by others in the sIL-6R and/or IL-6 moieties of the chimeric protein, such that modifications of this kind do not substantially change the biological activity of the chimeric protein analog with respect to the chimeric protein itself and in which the sIL-6R moiety of such analogs essentially retains the naturally occurring structure (before processing - see Fig. 3) of a signal peptide, Ig-like domain, cytokine receptor N-domain, cytokine receptor C-domain, and receptor pre-membrane domain. Likewise, such chimeric protein analogs should retain essentially the naturally-occurring mature form of the IL-6 moiety. The various analogs may differ most from each other and from the basic chimeric protein molecule (that with essentially only naturally-occurring sIL-6R and IL-6 sequences) at the site of the linker peptide which joins the sIL-6R and IL-6 moieties in the chimeric protein. Such a linker may be up to about 30 amino acids in length, and serves to separate the sIL-6R and IL-6 moieties from each other in the chimeric protein. As regards such a linker, care should be taken to choose its sequence (and hence also to test biologically in appropriate standard assays each such analog) such that it will, for example, not result in incorrect folding of the chimeric protein which may render it inactive, or it will not result in rendering the chimeric protein analog an immunogenic protein which will elicit antibodies against it in a patient to be treated therewith with the result that such an analog will be ineffective at least as a medium- or long- term medicament.

As regards the above analogs of the chimeric protein of the invention, these analogs are those in which one or more and up to about 30 of the amino

acid residues of the basic chimeric protein of the invention are replaced by different amino acid residues, or are deleted, or one or more amino acid residues are added to the original sequence of chimeric protein of the invention (that with essentially only the naturally-occurring sIL-6R and IL-6 sequences) without changing considerably the activity of the resulting products as compared with basic chimeric protein of the invention. These analogs are prepared by known synthesis and/or by site-directed mutagenesis techniques, or any other known technique suitable therefor.

Any such analog preferably has a sequence of amino acids sufficiently duplicative of that of the basic sIL-6R/IL-6 chimera such as to have substantially similar activity thereto. Thus, it can be determined whether any given analog has substantially the same activity as the basic chimeric protein of the invention by means of routine experimentation comprising subjecting such an analog to the biological activity tests set forth in Examples 2-4 below.

Analogues of the chimeric protein which can be used in accordance with the present invention, or nucleic acid coding therefor, include a finite set of substantially corresponding sequences as substitution peptides or polynucleotides which can be routinely obtained by one of ordinary skill in the art, without undue experimentation, based on the teachings and guidance presented herein. For a detailed description of protein chemistry and structure, see Schulz, G.E. et al., *Principles of Protein Structure*, Springer-Verlag, New York, 1978; and Creighton, T.E., *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, 1983, which are hereby incorporated by reference. For a presentation of nucleotide sequence substitutions, such as codon preferences, see Ausubel et al, *supra*, at §§ A.1.1-A.1.24, and Sambrook et al, Current Protocols in Molecular Biology, Interscience N.Y. §§6.3 and 6.4 (1987, 1992), at Appendices C and D.

Preferred changes for analogs in accordance with the present invention are what are known as "conservative" substitutions. Conservative amino acid substitutions of those in the chimeric protein having essentially the naturally-occurring sIL-6R and IL-6 sequences, may include synonymous amino acids within a group which have sufficiently similar physicochemical properties that substitution between members of the group will preserve the biological function of the molecule, Grantham, Science, Vol. 185, pp. 862-864 (1974). It is clear that insertions and deletions of amino acids may also be made in the above-defined sequences without altering their function, particularly if the insertions or deletions only involve a few amino acids, e.g., under thirty, and preferably under ten, and do not remove or displace amino acids which are critical to a functional conformation, e.g., cysteine residues, Anfinsen, "Principles That Govern The Folding of Protein Chains", Science, Vol. 181, pp. 223-230 (1973). Analogs produced by such deletions and/or insertions come within the purview of the present invention.

Preferably, the synonymous amino acid groups are those defined in Table I. More preferably, the synonymous amino acid groups are those defined in Table II; and most preferably the synonymous amino acid groups are those defined in Table III.

TABLE I Preferred Groups of Synonymous Amino Acids

Amino Acid	Synonymous Group
Ser	Ser, Thr, Gly, Asn
Arg	Arg, Gln, Lys, Glu, His
Leu	Ile, Phe, Tyr, Met, Val, Leu
Pro	Gly, Ala, Thr, Pro
Thr	Pro, Ser, Ala, Gly, His, Gln, Thr
Ala	Gly, Thr, Pro, Ala
Val	Met, Tyr, Phe, Ile, Leu, Val
Gly	Ala, Thr, Pro, Ser, Gly
Ile	Met, Tyr, Phe, Val, Leu, Ile
Phe	Trp, Met, Tyr, Ile, Val, Leu, Phe
Tyr	Trp, Met, Phe, Ile, Val, Leu, Tyr
Cys	Ser, Thr, Cys
His	Glu, Lys, Gln, Thr, Arg, His
Gln	Glu, Lys, Asn, His, Thr, Arg, Gln
Asn	Gln, Asp, Ser, Asn
Lys	Glu, Gln, His, Arg, Lys
Asp	Glu, Asn, Asp
Glu	Asp, Lys, Asn, Gln, His, Arg, Glu
Met	Phe, Ile, Val, Leu, Met
Trp	Trp

TABLE II More Preferred Groups of Synonymous Amino Acids

Amino Acid	Synonymous Group
Ser	Ser
Arg	His, Lys, Arg
Leu	Leu, Ile, Phe, Met
Pro	Ala, Pro
Thr	Thr
Ala	Pro, Ala
Val	Val, Met, Ile
Gly	Gly
Ile	Ile, Met, Phe, Val, Leu
Phe	Met, Tyr, Ile, Leu, Phe
Tyr	Phe, Tyr
Cys	Cys, Ser
His	His, Gln, Arg
Gln	Glu, Gln, His
Asn	Asp, Asn
Lys	Lys, Arg
Asp	Asp, Asn
Glu	Glu, Gln
Met	Met, Phe, Ile, Val, Leu
Trp	Trp

TABLE III Most Preferred Groups of Synonymous Amino Acids

Amino Acid	Synonymous Group
Ser	Ser
Arg	Arg
Leu	Leu, Ile, Met
Pro	Pro
Thr	Thr
Ala	Ala
Val	Val
Gly	Gly
Ile	Ile, Met, Leu
Phe	Phe
Tyr	Tyr
Cys	Cys, Ser
His	His
Gln	Gln
Asn	Asn
Lys	Lys
Asp	Asp
Glu	Glu
Met	Met, Ile, Leu
Trp	Met

Examples of production of amino acid substitutions in proteins which can be used for obtaining analogs of the chimeric protein for use in the present invention include any known method steps, such as presented in US patents RE 33,653, 4,959,314, 4,588,585 and 4,737,462, to Mark et al; 5,116,943 to Koths

et al., 4,965,195 to Namen et al; 4,879,111 to Chong et al; and 5,017,691 to Lee et al; and lysine substituted proteins presented in US patent No. 4,904,584 (Shaw et al).

In another preferred embodiment of the present invention, any analog of the chimeric protein for use in the present invention has an amino acid sequence essentially corresponding to that of the above noted basic chimeric protein of the invention. The term "essentially corresponding to" is intended to comprehend analogs with minor changes to the sequence of the basic chimeric protein which do not affect the basic characteristics thereof, particularly insofar as its ability to inhibit cancer cell proliferation or promote bone marrow transplantations, for example, is concerned. The type of changes which are generally considered to fall within the "essentially corresponding to" language are those which would result from conventional mutagenesis techniques of the DNA encoding the chimeric protein of the invention, resulting in a few minor modifications, and screening for the desired activity in the manner discussed above.

Analogues in accordance with the present invention include those encoded by a nucleic acid, such as DNA or RNA, which hybridizes to DNA or RNA under stringent conditions and which encodes a chimeric protein in accordance with the present invention, comprising essentially all of the naturally-occurring sequences encoding sIL-6R and IL-6. For example, such a hybridizing DNA or RNA maybe one encoding the same protein of the invention having, for example, the sequence set forth in Fig. 3, but which nucleotide differs in its nucleotide sequence from the naturally-derived nucleotide sequence by virtue of the degeneracy of the genetic code, i.e., a somewhat different nucleic acid sequence may still code for the same amino acid sequence, due to this degeneracy. Further, as also noted above, the amount of amino acid changes (deletions, additions, substitutions) is limited to up to about 30 amino acids, such that even with the maximum amount of changes, analogues in accordance with the present invention

will be those which essentially retain the leader (before processing), Ig-like domain, cytokine receptor N- and C- domains and receptor pre-membrane region in the sIL-6R moiety and essentially all of the IL-6 moiety. Such nucleic acid would be a prime candidate to determine whether it encodes a polypeptide which retains the functional activity of the chimeric protein of the present invention. The term "stringent conditions" refers to hybridization and subsequent washing conditions which those of ordinary skill in the art conventionally refer to as "stringent". See Ausubel et al., Current Protocols in Molecular Biology, supra, Interscience, N.Y., para. 6.3 and 6.4 (1987, 1992), and Sambrook et al., supra. Without limitation, examples of stringent conditions include washing conditions 12-20°C below the calculated T_m of the hybrid under study in, e.g. 2 x SSC and 0.5% SDS for 5 minutes, 2 x SSC and 0.1% SDS for 15 minutes; 0.1 x SSC and 0.5% SDS at 37°C for 30-60 minutes and then a 0.1 x SSC and 0.5% SDS at 68°C for 30-60 minutes. Those of ordinary skill in this art understand that stringency conditions also depend on the length of the DNA sequences, oligonucleotide probes (such as 10-40 bases) or mixed oligonucleotide probes. If mixed probes are used, it is preferable to use tetramethyl ammonium chloride (TMAC) instead of SSC. See Ausubel, supra.

The term "salts" herein refers to both salts of carboxyl groups and to acid addition salts of amino groups of the chimeric protein of the invention or analogs thereof. Salts of a carboxyl group may be formed by means known in the art and include inorganic salts, for example, sodium, calcium, ammonium, ferric or zinc salts, and the like, and salts with organic bases as those formed, for example, with amines, such as triethanolamine, arginine or lysine, piperidine, procaine and the like. Acid addition salts include, for example, salts with mineral acids such as, for example, hydrochloric acid or sulfuric acid, and salts with organic acids such as, for example, acetic acid or oxalic acid. Of course, any such salts must

have substantially similar activity to the chimeric protein of the invention or its analogs.

The present invention also concerns DNA sequences encoding the above chimeric protein of the invention and its analogs, as well as DNA vectors carrying such DNA sequences for expression in suitable mammalian, preferably human, cells. An embodiment of a vector of the invention is a plasmid pcDNA sIL-6R/IL-6 comprising the pcDNA3 vector (Invitrogen) containing the sIL-6R/IL-6 fused sequences under the control of a cytomegalovirus (CMV) promoter.

The present invention also concerns transformed mammalian, preferably human, cells capable of expressing the above vectors of the present invention. An embodiment of such transformed cells are human embryonal kidney cells 293 (HEK 293, ATCC CRL 1573) transfected by pcDNA sIL-6R/IL-6 which secrete the fused sIL-6R/IL-6 chimeric as a 85 kDa glycoprotein.

A further embodiment is plasmid pcDNA sIL-6R/L/IL-6 which differ from the above pcDNA sIL-6R/IL-6 by insertion in the EcoRI site of short linkers encoding 10 additional aminoacids. A number of different sequences, of various lengths, can be introduced to optimize the distance between sIL-6R and IL-6.

The present invention further concerns a method for producing and purifying the chimeric protein of the invention or its analogs which comprises growing the above transformed cells under conditions suitable for expression and secretion of the chimeric protein product into the culture medium and then purifying the secreted protein by immunoaffinity chromatography using anti-sIL-6R monoclonal antibodies 34.4 as noted in Example 2 below.

The invention also concerns a pharmaceutical composition comprising as active ingredient an sIL-6R/IL-6 chimera or analogs thereof or mixtures thereof or salts thereof and a pharmaceutically acceptable carrier, diluent or excipient. An embodiment of the pharmaceutical composition of the invention include a pharmaceutical composition for enhanced IL-6 type action, for the treatment of

cancers, for bone marrow transplantation, for increase of hematopoiesis, in particular thrombopoiesis, for treatment of neurological conditions and other applications of IL-6 or related cytokines.

The pharmaceutical compositions of the invention are prepared for administration by mixing the chimeric protein, or its analogs with physiologically acceptable carriers, and/or stabilizers and/or excipients, and prepared in dosage form, e.g., by lyophilization in dosage vials. The method of administration can be via any of the accepted modes of administration for similar agents and will depend on the condition to be treated, e.g., intravenously, intramuscularly, subcutaneously, by local injection or topical application, or continuously by infusion, etc. The amount of active compound to be administered will depend on the route of administration, the disease to be treated and the condition of the patient. Local injection, for instance, will require a lower amount of the protein on a body weight basis than will intravenous infusion.

The present invention also concerns uses of the chimeric protein of the invention or its analogs or mixtures thereof for the treatment of cancers, for bone marrow transplantations, for increasing hematopoiesis, especially thrombopoiesis, for treatment of neurological conditions and for use in other applications of IL-6 or related cytokines. Likewise, the present invention also concerns the chimeric protein or analogs thereof or mixtures thereof for use in the preparation of medicaments for treating the above-mentioned ailments or for use in the above noted indications.

The present invention will now be described in more detail in the following non-limiting Examples and the accompanying drawings.

Example 1: Construction of the sIL-6R δ Val/IL-6 chimera expression vector

In Figure 1, there is shown a schematic flow-diagram of the steps taken to construct the expression vector carrying the sequence coding for the sIL-6R δ Val/IL-6 chimeric protein, inclusive of all the various starting and intermediary vectors, various reagents and reaction steps. This construction procedure was essentially using techniques well known in the art for constructing expression vectors of choice (see, for example, Sambrook et al., 1989). The procedure was, briefly, as follows :

A library of cDNAs from human breast carcinoma T47D cells was cloned in the lamda (λ) gt11 bacteriophage and screened with oligonucleotide probes derived from the IL-6R sequence of Yamasaki et al (1988). One λ gt11 cDNA clone was isolated which had the entire human IL-6R coding sequence. The insert was excised from λ gt11 by EcoRI and cloned in the Multiple Cloning Site (MCS) of the E.coli phagemid Blue Script pBS/SK (Stratagene Cloning Systems, LaJolla, California). This plasmid pBS/SK-IL-6R (Figure 1) was cut by EcoRI which was then blunt-ended and recut with EcoRV to isolate the 5' fragment of IL-6R of 959 base pairs (bp) ending at the EcoRV site of IL-6R (coordinate 1203). This fragment extracted from an agarose gel electrophoresis was cloned in a new pBS/SK vector opened at the EcoRV of the MCS (pBS/SK-sIL6R-RV in Figure 1).

The above noted, previously obtained pBS/SK-IL-6R DNA was subjected to Polymerase Chain Reaction (PCR) to amplify a 368 bp fragment between the forward primer 1137-1156 and the reverse primer 1505-1488. The reverse primer was synthesized with an EcoRI site immediately following the codon for Valine-356 of the IL-6R (see Figure 1), since this Valine residue was previously determined to be the carboxy-terminal amino acid of the natural form of the soluble sIL-6R excreted in human urine (Novick et al, 1990; Oh et al, 1996; co-owned U.S. Pat. No. 5,216,128 and EP Pat. No. EP 413908 B1). The PCR product was cut by EcoRV and by EcoRI and ligated into pBS/SK-sIL6R-RV between the EcoRV site

of IL-6R and the EcoRI site of the MCS (Figure 1). The resulting plasmid pBS-sIL6R- δ Val-RI was then shortened to remove 5' untranslated sequences by ligation of the HindIII site of MCS with the NcoI site at base pair 410 of IL-6R (both sites being first blunt-ended), to yield pBS-sIL6R- δ Val-RI-NcoI (Figure 1).

The IL-6 sequence was derived from plasmid pKK β 2-7 which, as previously described (Chen et al, 1988), was constructed by insertion of the BstNI-cut IFN- β 2/IL-6 cDNA (Zilberstein et al, 1986) into the EcoRI site of the E.coli expression vector pKK223-3 (Pharmacia, Uppsala, Sweden) using a synthetic oligonucleotide with an EcoRI site followed by a Methionine codon and the codon for Proline-29 of IL-6 and ending at a BstNI (EcoRII) site. The IL-6 cDNA insert of pKK β 2-7 ends 7 base pairs after the termination codon a NlaIV site and is followed 11 bp later by the HindIII site of the pKK223-3 vector (Figure 1). The pKK β 2-7 DNA was cut with HindIII, blunt-ended and recut with EcoRI and the IL-6 cDNA inserted into pBS-sIL6R- δ Val-RI-NcoI so as to fuse the mature sequence of IL-6 (starting at Proline-29) immediately after Valine-356 of the IL-6R and separated by only 3 codons (Glu-Phe-Met). The resulting plasmid pBS/SK-sIL6R/IL-6 (Figure 1) was then recut at the SalI and NotI sites of its MCS and the insert was cloned into the EcoRV site of pcDNA3 (Invitrogen Corporation, San Diego, California). The resulting plasmid pcDNA3-sIL6R/IL-6 (Figure 1) contains the insert downstream of the strong cytomegalovirus (CMV) promoter and followed by a polyadenylation site insuring efficient transcription of the sIL6R δ Val/IL6 chimera. The conservation of the 5' end of the sIL-6R in the chimera ensures that upon expression in mammalian cells the signal peptide function and processing of the N-terminus of the chimeric protein will be as in the natural sIL-6R.

As indicated above, an advantageous characteristic of the sIL6R δ Val/IL6 construct is that it is essentially the fusion of the natural form of sIL-6R and of the natural form of IL-6 as they exist in the human body, and without extraneous polypeptide sequences. However, the conservation of the EcoRI site in the

sIL6R δ Val/IL6 construct (Figure 1) allows to easily introduce linker polypeptide segments between the sIL-6R and the IL-6 moieties. One such construct with the 13-amino acid linker sequence Glu-Phe-Gly-Ala-Gly-Leu-Val-Leu-Gly-Gly-Gln-Phe-Met introduced between Val-356 of sIL-6R and Pro-29 of IL-6, was also constructed (sIL6R δ Val/L/IL6).

Example 2: Expression of the sIL-6R δ Val/IL-6 chimera in human cells.

Using essentially standard techniques of mammalian cell culture, cell transfection and analysis of the transfected cells for expression of the newly introduced DNA sequence to be expressed (for procedures, see, for example, Sambrook et al., 1989), the above plasmid construct (Example 1) was used to transfect human cells and its expression therein was assessed. Briefly, the following procedures were employed :

Human HEK 293 cells (ATCC CRL 1573, transformed primary human embryonal kidney cells) were transfected with the plasmic construct pCDNA3-sIL6R/IL6 DNA (set forth in Example 1 above). Log phase cultures of HEK 293 were trypsinized and seeded in 9cm Nunc plates (2.5×10^6 cells/plate). One day later, transfection was carried out with 10 μ g pCDNA3-sIL6R/IL6 DNA by the CaPO₄ precipitation procedure (Sambrook et al, 1989) and 1 hour later the medium changed to DMEM-10% FCS and the culture continued for an additional 16 hours. The secreted proteins were then collected in DMEM-2% FCS for two consecutive periods of 48 hours. Debris was removed by centrifugation at 1,000 rpm for 10 minutes and the supernatant tested by an ELISA for sIL-6R using polyclonal rabbit anti-sIL-6R and mouse McAB 17.6 (Novick et al, 1991). A concentration of 1.2 μ g/ml sIL-6R -equivalents was found, indicative of very efficient expression of the chimeric sIL-6R/IL-6 protein in the transfected human cells.

Immunopurification of the secreted chimeric protein (sIL-6R/IL-6) was carried out with Monoclonal Antibody 34.4 specific to an epitope in the

extracellular domain of human sIL-6R (Novick et al, 1991; Halimi et al, 1995). The 34.4 hybridoma cells were grown in the peritoneal cavity of mice and the immunoglobulin (Ig) fraction was obtained from the ascitis fluid by ammonium sulfate precipitation. Affigel-10 (Bio-Rad Labs, Richmond, California) was used to immobilize McAB 34.4 (15 mg Ig coupled to 1 ml Affigel-10). The supernatants containing the secreted proteins from the HEK 293 cells transfected by pCDNA3-sIL6R/IL6 were adsorbed on columns of McAB 34.4 (0.3 ml column for 15 ml supernatant). After washing with PBS, the bound proteins were eluted by 25 mM citric acid pH 2.5, then immediately neutralized by 1 M Hepes buffer pH 8.5 and dialyzed overnight (about 8-12 hrs) against PBS.

Analysis of the immunopurified protein by polyacrylamide gel electrophoresis in SDS showed a unique protein band stained by Coomassie blue (Figure 2). The molecular weight of the protein was 85 kilodaltons as expected from the fusion of the glycosylated forms of sIL-6R δ Val (60 kDa as shown in Oh et al, 1996) and glycosylated IL-6 (23-26 kDa as shown in Zilberstein et al, 1986). The aminoacid sequence of the sIL6R/IL6 is 543 aminoacids, which after processing of the signal peptides would predict a protein of 524 aminoacids or about 58 kDa (Figure 3). The much larger size of the sIL6R/IL6 chimera produced from the recombinant DNA in human cells indicates that glycosylation accounts for a sizable portion of the molecule.

Example 3: The sIL6R/IL6 chimera arrests growth and induces differentiation of metastatic melanoma cells.

The F10.9 clone derived from B16 melanoma cells forms highly metastatic tumors in C57Black/6 mice which kill the animals from pulmonary metastases within 2-3 months (Katz et al, 1995). Addition of the sIL6R/IL6 chimeric protein to F10.9 cells culture produces a profound morphological change in the cells and an arrest in their growth (Figure 4). The F10.9 cells treated by the chimera become

elongated, with protruding dendritic extensions, resembling the spindloid differentiation of embryonic melanocytes.

The growth of the cells was quantitated 4 days after seeding 3×10^3 cells in wells of a 96-well microplate in 0.2 ml RPMI 1640 medium with 10% FCS. The cells were fixed in 12.5% glutaraldehyde for 30 minutes, washed in water and stained with 0.1% crystal violet for 30 minutes. After thorough washing and drying, the stain was extracted by 10% acetic acid and the optical density determined at 540 nm. The chimera produced a dose-dependent inhibition of growth with a complete growth inhibition at concentrations as low as 10 ng/ml of the chimeric (p85) protein (Figure 5). Both chimeric proteins sIL6R δ Val/IL6 and sIL6R δ Val/L/IL6 (chimera with the longer linker between the sIL-6R and IL-6 moieties, see Example 1) were similarly active. This result also serves to show that the linker peptide between the sIL-6R and IL-6 moieties in the chimera, is not essential for the activity of the chimera as the above sIL-6R δ Val/IL-6 chimera has only a very short 3 amino acid linker while the above sIL-6R δ Val/L/IL-6 has a longer 13 amino acid linker peptide, but both have essentially the same activity in inhibiting the growth of the metastatic cells. In contrast, neither IL-6 alone, nor the sIL-6R δ Val alone inhibit the growth of these melanoma cells (Figure 6) demonstrating the unique activity of the sIL-6R/IL-6 (p85) chimeric protein. To obtain a similar effect, a mixture of 200-400 ng/ml IL-6 and 125 ng/ml sIL-6R δ Val is required (Figure 6). When calculated in molar concentrations, the maximal inhibition of F10.9 cells required 7.5 nM IL-6 and 2 nM sIL-6R δ Val versus only 0.12 nM of the sIL-6R/IL-6 chimera.

The growth inhibitory activity of the p85 sIL6R/IL6 chimeric protein was followed during the immunopurification on McAB 34.4 columns (see Example 2). The pattern of activity corresponded to the intensity of the p85 band seen in the different fractions of the SDS polyacrylamide gel electrophoresis in Figure 2.

Example 4: The sIL6R/IL6 Chimera is essential for engraftment of human bone marrow transplanted cells

Engraftment of hematopoietic stem cells from human bone marrow can be studied after transplantation into severe combined immunodeficient (SCID) mice (Vormoor et al, 1994). SCID-NOD mice were subjected to sublethal irradiation and injected in the tail vein with 3×10^5 human CD34⁺ bone marrow cells. Prior to injection, the purified CD34⁺ cells were maintained for 3 days in liquid cultures with different combinations of cytokines. After one month, the mice were sacrificed and bones were taken to collect the bone marrow cells. The engraftment of human cells in the SCID-NOD recipient mice was evaluated by Southern blot hybridization to human repetitive DNA.

Stem-cell factor (SCF, steel factor or ckit-ligand) and Flt3-ligand (flt3/flk2 tyrosine kinase receptor ligand) have been found important for survival and proliferation of the most primitive pluripotential hematopoietic stem cells capable of long-term engraftment in recipient bone marrow (McKenna et al, 1995). As seen in Figure 7, these two factors by themselves were insufficient to promote the engraftment of human cells in the bone marrow of the SCID-NOD recipient mice. Addition of the sIL6R/IL6 chimeric protein was required for engraftment to be detected at significant levels. At 100 ng/ml, the sIL6R/IL6 chimera was much more active than the isolated IL-6 (50-200 ng/ml) and sIL-6R (125-1250 ng/ml) (Figure 7). The requirement for the sIL6R/IL6 chimera indicates that this protein is essential for the survival and proliferation of the non-committed pluripotential hematopoietic stem cells which can home into and repopulate the bone marrow environment, indicating that this protein may be useful in bone marrow transplantation clinical protocols.

This is the first demonstration that the sIL-6R/IL-6 chimera has the following at least two newly found activities :

(i) When added together with both of the factors SCF and Flt3- ligand to human hematopoietic primitive progenital cells, it promotes their survival and proliferation; and

(ii) It is active (and apparently essential) in an *in vivo* model of a human bone marrow transplantation in immuno-deficient mice.

References

- Chen L, Mory Y, Zilberstein A and Revel M. Growth inhibition of human breast carcinoma and leukemia/lymphoma cell lines by recombinant interferon-beta 2/IL-6. *Proc. Natl. Acad. Sci. USA*, 85: 8037-8041, 1988.
- Chernajovsky Y, Mory Y, Chen L, Marks Z, Novick D, Rubinstein M and Revel M. Efficient constitutive production of human fibroblast interferon by hamster cells transformed with the IFN- β 1 gene fused to an SV40 early promoter. *DNA*, 3: 297-308, 1984.
- Fischer M, Goldschmitt J, Peschel C, Brakenhoff JPG, Kallen K-J, Wollmer A, Grotzinger J and Rose-John S. A bioactive designer cytokine for human hematopoietic progenitor cell expansion. *Nature Biotechnology* 15: 142-145, 1997.
- Ganapathi MK, Weizer AK, Borsellino S, Bukowski RM, Ganapathi S, Rice T, Casey G and Kawamura K. Resistance to Interleukin-6 in human Non-small cell lung carcinoma cell lines: Role of receptor components. *Cell Growth and Differentiation*, 7: 923-929, 1996.
- Halimi H, Eisenstein M, Oh J, Revel M and Chebath J. Epitope peptides from interleukin-6 receptor which inhibit the growth of human myeloma cells. *Eur. Cytokine Netw.*, 6: 135-143, 1995.
- Hirano T, Yasukawa K, Harada H, Taga T, Watanabe Y, Matsuda T, Kashimura S, Nakajima K, Koyama K, Iwamatsu K, Tsunasawa S, Sakiyama F, Matsui H, Takahara Y, Taniguchi T and Kishimoto T. Complementary DNA for a novel interleukin (BSF-2) that induces B lymphocytes to produce immunoglobulins. *Nature*, 234: 73-76, 1986.
- Hirano T, Matsuda T and Nakajima K. Signal transduction through gp130 that is shared among the receptors for the interleukin 6 related cytokine subfamily. *Stem cells*, 12:262-277, 1994.

- Holloway CJ. Applications of recombinant DNA technology in the production of glycosylated recombinant human granulocyte colony stimulating factor. *Eur. J. Cancer*, 30: S2-6, 1994.
- Kahn MA and De Vellis J. Regulation of an oligodendrocyte progenitor cell line by the interleukin-6 family of cytokines. *Glia*, 12: 87-98, 1994.
- Katz A, Shulman LM, Porgador A, Revel M, Feldman M and Eisenbach L. Abrogation of B16 melanoma metastases by long-term low-dose Interleukin-6 therapy. *J. Immunother.* 13: 98-109, 1993.
- Mackiewicz A, Wiznerowicz M, Roeb E, Nowak J, Pawlowski T, Baumann H, Heinrich P and Rose-John S. Interleukin-6-type cytokines and their receptors for gene therapy of melanoma. *Ann. New York Acad. Sci.*, 762: 361-374, 1995.
- McKenna HJ, de Vries P, Brasel K, Lyman SD and Williams DE. Effect of flt3 ligand on the ex vivo expansion of human CD34+ hematopoietic progenitor cells. *Blood* 86: 3413-3420, 1995.
- Murakami M, Hibi M, Nakagawa N, Nagakawa T, Yasukawa K, Yamanishi K, Taga T and Kishimoto T. IL-6 induced homodimerization of gp130 and associated activation of a tyrosine kinase. *Science*, 260: 1808-1810, 1993.
- Novick D, Englemann H, Wallach D, Leitner O, Revel M and Rubinstein M. Purification of soluble cytokine receptors from normal urine by ligand-affinity and immunoaffinity chromatography. *J. Chromatogr.*, 510: 331-337, 1990.
- Novick D, Englemann H, Revel M, Leitner O and Rubinstein M. Monoclonal antibodies to the soluble IL-6 receptor: affinity purification, ELISA and inhibition of ligand binding. *Hybridoma*, 10: 137-146, 1991.
- Novick D, Shulman LM, Chen L and Revel M. Enhancement of interleukin-6 cytostatic effect on human breast carcinoma cells by soluble IL-6 receptor from urine and reversion by monoclonal antibodies. *Cytokine*, 4: 6-11, 1992.

Oh J-W, Revel M and Chebath J. A soluble interleukin-6 receptor isolated from conditioned medium of human breast cancer cells is encoded by a differentially spliced mRNA. *Cytokine*, 8: 401-409, 1996.

Oh J-W. Expression of recombinant soluble human interleukin-6 receptors and analysis of their functions. Ph.D. Thesis Weizmann Institute of Science (Revel M, supervisor), 1997.

Paonessa G, Graziani R, DeSerio A, Savino R, Ciapponi L, Lahmm A, Salvati AL, Toniatti C and Ciliberto G. Two distinct and independent sites on IL-6 trigger gp130 dimer formation and signalling. *EMBO J.*, 14: 1942-1951, 1995.

Revel M. Host defense against infections and inflammations: Role of the multifunctional IL-6/IFN- β 2 cytokine. *Experientia* 45: 549-557, 1989.

Sambrook J, Fritsch EF and Maniatis T. *Molecular cloning: A laboratory manual*. Cold Spring Harbor Press, 1989.

Sui X, Tsuji K, Tanaka R, Tajima S, Muraoka K, Ebihara Y, Ikebuchi K, Yasukawa K, Taga T, Kishimoto T and Nakahata T. Gp130 and c-kit signalings synergize for ex vivo expansion of human primitive hemopoietic progenitor cells. *Proc. Natl. Acad. Sci. USA* 92: 2859-2863, 1995.

Taga T, Hibi M, Hirata Y, Yamasaki K, Yasukawa K, Matsuda T, Hirano T and Kishimoto T. Interleukin-6 triggers the association of its receptor with a possible signal transducer gp130. *Cell*, 58: 573-581, 1989.

Vormoor J, Lapidot T, Pflumio F, Risdon G, Patterson B, Broxmeyer HE and Dick JE. SCID mice as an in vivo model of human cord blood hematopoiesis. *Blood cells* 20: 316-320, 1994.

Ward LD, Howlett GJ, Discolo G, Yasukawa K, Hammacher A, Moritz RL and Simpson RJ. High affinity interleukin-6 receptor is a hexameric complex consisting of two molecules each of interleukin-6, interleukin-6 receptor and gp130. *J. Biol. Chem.*, 269: 23286-23289, 1994.

Yamasaki K, Taga T, Hirata Y, Yawata H, Kawanishi Y, Seed B, Taniguchi T, Hirano T and Kishimoto T. Cloning and expression of the human Interleukin-6 (BSF-2/ Interferon beta-2) receptor. *Science*, 241: 825-828, 1988.

Zilberstein A, Ruggieri R, Korn HJ and Revel M. Structure and expression of of cDNA and genes for human interferon-beta-2, a distinct species inducible by growth-stimulatory cytokines. *EMBO J.*, 5: 2529-2537, 1986.

CLAIMS

1. A chimeric glycosylated soluble interleukin-6 receptor (sIL-6R)-interleukin-6 (IL-6) protein (sIL-6R/IL-6) and biologically active analogs thereof, comprising a fusion protein product between essentially all of the naturally occurring form of sIL-6R and essentially all of the naturally occurring form of IL-6, said sIL-6R/IL-6 and analogs thereof being glycosylated in a similar fashion to the glycosylation of naturally occurring sIL-6R and IL-6.
2. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to claim 1, wherein said sIL-6R is fused to IL-6 via a peptide linker molecule.
3. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof, according to claim 2, wherein said linker is a very short, non-immunogenic linker of about 3-4 amino acid residues.
4. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to claim 3, wherein said linker is a tripeptide of the sequence E-F-M (Glu-Phe-Met).
5. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to claim 2, wherein said linker is a peptide of 13 amino acid residues of sequence E-F-G-A-G-L-V-L-G-G-Q-F-M (Glu-Phe-Gly-Ala-Gly-Leu-Val-Leu-Gly-Gly-Gln-Phe-Met).
6. A chimeric sIL-6R/IL-6 protein according to any one of claims 1-4, being the herein designated sIL6R δ Val/IL6 having a tripeptide linker of sequence E-F-M

between the C-terminal Val-356 of sIL-6R and the N-terminal Pro-29 of IL-6R, said chimeric protein having the sequence set forth in Fig. 3.

7. A chimeric sIL-6R/IL-6 protein according to any one of claims 1, 2, and 5, being the herein designated sIL6R δ Val/L/IL6 having a 13 amino acid peptide linker of sequence E-F-G-A-G-L-V-L-G-G-Q-F-M between the C-terminal Val-356 of sIL-6R and the N-terminal Pro-29 of IL-6R, said chimeric protein having the sequence set forth in Fig. 3 wherein the tripeptide of sequence E-F-m between positions 357-359 of Fig. 3 is replaced by said 13 amino acid peptide sequence.

8. A chimeric sIL-6R/IL-6 protein according to any one of claims 1-7, wherein said protein is produced in mammalian cells in a fully processed form.

9. A chimeric sIL-6R/IL-6 protein according to claim 8, wherein said protein is produced in human cells.

10. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to any one of claims 1-9, wherein said chimeric protein and analogs are characterized by being capable of inhibiting the growth of highly malignant cancer cells.

11. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to claim 10, wherein said chimeric protein and analogs are characterized by being capable of inhibiting the growth of highly malignant melanoma cells.

12. A chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to any one of claims 1-9, wherein said chimeric protein and analogs are

characterized by being capable of eliciting the *in vivo* engraftment of human hematopoietic cells in bone marrow transplantations.

13. A DNA sequence encoding a chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to any one of claims 1-12.

14. A DNA vector comprising a DNA sequence encoding a chimeric sIL-6R/IL-6 protein and biologically active analogs thereof according to any one of claims 1-12, said vector being suitable for expression of said chimeric protein in mammalian cells.

15. A DNA vector according to claim 14, wherein said vector is suitable for expression of said chimeric protein in human cells.

16. A DNA vector according to claim 14 or 15, wherein when said vector is expressed in mammalian or human cells, the expressed chimeric protein has a sequence that permits full processing of the chimeric protein by the mammalian or human cell and secretion of the fully processed chimeric protein from the cells into the culture medium in which said cells are grown.

17. A DNA vector according to any one of claims 14-16, wherein said vector is the herein designated plasmid pcDNAsIL-6R/IL-6 comprising a pcDNA3 vector containing the DNA sequence encoding the chimeric sIL-6R/IL-6 protein under the control of a cytomegalovirus (CMV) promoter.

18. A DNA vector according to any one of claims 14-16, wherein said vector is the herein designated plasmid pcDNA sIL-6R/L/IL-6 comprising a pcDNA3 vector containing the DNA sequence encoding the chimeric sIL-6R/IL-6 protein under the

control of a cytomegalovirus (CMV) promoter, and wherein in said DNA sequence encoding said chimeric sIL-6R/IL-6 protein there is inserted a linker sequence encoding a peptide linker at the EcoRI site placed between the sequence encoding the sIL-6R part and the sequence encoding the IL-6 part of the protein.

19. Transformed mammalian cells containing a DNA vector according to any one of claims 14-18 which are capable of expressing the sIL-6R/IL-6 chimeric protein sequence carried by said vector and of fully processing the expressed protein and secreting it into the culture medium in which said cells are grown.

20. Transformed cells according to claim 19 wherein in said cell is the herein described human embryonal kidney cells 293 (HEK293) transfected by the pcDNA sIL-6R/IL-6 vector, said cells being capable of expressing the sIL-6R/IL-6 chimeric protein, fully processing said protein and secreting said protein into the culture medium in which said cells are grown in the form of an about 85 kDa glycoprotein.

21. A method for producing a chimeric protein or biologically active analogs thereof according to any one of claims 1-12, comprising growing transformed cells according to claim 19 or 20 under conditions suitable for expression, processing and secretion of said protein or analogs into the culture medium in which said cells are grown; and purifying said protein or analogs from said culture medium.

22. A method according to claim 21, wherein the purification is carried out by immunoaffinity chromatography using monoclonal antibodies specific for sIL-6R.

23. The use of a chimeric sIL-6R/IL-6 protein or analogs according to any one of claims 1-12, salts of any one thereof, and mixtures thereof, as an inhibitor of cancer cells.

24. The use of a chimeric protein or analog according to claim 23, as an inhibitor of highly malignant melanoma cells.

25. The use of a chimeric sIL-6R/IL-6 protein or analogs according to any one of claims 1-9 and 12, salts of any one thereof, and mixtures thereof, as an active ingredient for eliciting engraftment of human hematopoietic cells in bone marrow transplantation.

26. The use of a chimeric sIL-6R/IL-6 protein or analogs according to any one of claims 1-9 and 12, salts of any one thereof, and mixtures thereof, as an active ingredient for increasing hematopoiesis, for treating neurological conditions, or for other applications in which IL-6 or sIL-6R are used.

27. A chimeric sIL-6R/IL-6 protein or analogs according to any one of claims 1-12, salts of any one thereof and mixtures thereof, for use in the preparation of a medicament for treating mammalian cancers by way of inhibition of mammalian cancer cells, or in the preparation of a medicament for enhancement of bone marrow transplantation by way of eliciting engraftment of human hematopoietic cells in bone marrow transplantation, or in the preparation of a medicament for increasing hematopoiesis, or in the preparation of a medicament for treating neurological disorders, or in the preparation of a medicament for other applications in which IL-6 or sIL-6R are used.

28. A pharmaceutical composition comprising as active ingredient a chimeric sIL-6R/IL-6 protein or analog thereof according to any one of claims 1-12, and a pharmaceutically acceptable carrier, diluent or excipient.

29. A pharmaceutical composition according to claim 28 for the treatment of cancers.

30. A pharmaceutical composition according to claim 28 for the enhancement of bone marrow transplantation.

31. A pharmaceutical composition according to claim 28 for the treatment of neurological disorders, or for increasing hematopoiesis or for other applications in which IL-6 or sIL-6R are used.

32. A method for treating cancers in mammals, or for enhancing bone marrow transplantations, or for treating neurological disorders, or for increasing hematopoiesis, or for other applications in which IL-6 or sIL-6R are used, comprising administering to a patient a pharmaceutical composition according to any one of claims 28-31 in a suitable dosage form and by a suitable route of administration.

For the Applicant



Henry Einav



1
2
3